

Chiral Symmetry Breaking for Domain Wall Fermions in Quenched Lattice QCD

Lingling Wu^{a*}

^aDepartment of Physics, Columbia University, New York, NY, 10027, USA

The domain wall fermion formulation exhibits full chiral symmetry for finite lattice spacing except for the effects of mixing between the domain walls. Close to the continuum limit these symmetry breaking effects should be described by a single residual mass. We determine this mass from the conservation law obeyed by the conserved axial current in quenched simulations with $\beta = 5.7$ and 6.0 and domain wall separations varying between 12 and 48 on $8^3 \times 32$ and $16^3 \times 32$ lattices. Using the resulting values for the residual mass we perform two complete and independent calculations of the pion decay constant. Good agreement is found between these two methods and with experiment.

1. INTRODUCTION

An important feature of both domain wall and overlap fermion formulations is that the chiral limit is disentangled from the continuum limit. For the case of the domain wall fermions, the chiral symmetry breaking is solely controlled by the amount of coupling between the domain walls[1]. To exploit the chiral properties of the domain wall fermions, it is essential to understand these chiral symmetry breaking effects.

We reported earlier[2,3] that for quenched simulations, although the quantity $m_\pi^2(m_f = 0)$ obtained from the pseudoscalar density correlator decreased as the separation between the domain walls L_s was increased, it did not appear to vanish in the $L_s \rightarrow \infty$ limit. Further investigation shows that the chiral limit of the pion mass is distorted by many factors. Besides the order a^2 effects and finite volume effects common to both dynamical and quenched simulations using domain wall fermions, the presence of topological near-zero modes in the quenched approximation significantly affects the determination of the pion mass[4,5]. These zero-mode effects can be

suppressed by going to a larger volume. Other non-linear behaviors caused by the absence of the fermion determinant in quenched simulations, such as “quenched chiral logs”, may also play an important role in the behavior of the pion mass as the chiral limit is approached. These factors make it difficult to determine the degree of chiral symmetry breaking from examination of the pion mass in the quenched chiral limit. The technique described in detail below quantifies the chiral symmetry breaking effects more precisely by a single residual mass measured from the extra term in the divergence of the axial current[6].

2. RESIDUAL CHIRAL SYMMETRY BREAKING

Close to the continuum, the effect of mixing between the domain walls produces chiral symmetry breaking terms with a coefficient m_{res} , the residual mass, in the low energy effective Lagrangian. This results in an effective quark mass of $m_{\text{eff}} = m_f + m_{\text{res}}$ entering all low momentum Green’s functions.

We can measure m_{res} starting from the divergence of the axial current, given by[6]

$$\Delta_\mu \mathcal{A}_\mu^a(x) = 2m_f J_5^a(x) + 2J_{5q}^a(x). \quad (1)$$

Compared with the corresponding continuum expression, the additional term from J_{5q}^a , referred as the “mid-point” contribution, should be equiva-

*This work was done in collaboration with T. Blum, P. Chen, N. Christ, C. Christian, C. Dawson, G. Fleming, A. Kaehler, X. Liao, G. Liu, C. Malureanu, R. Mawhinney, S. Ohta, G. Siebert, A. Soni, C. Sui, P. Vranas, M. Wingate, Y. Zhestkov(RIKEN-BNL-CU collaboration). This work was supported in part by the Department of Energy and the RIKEN BNL Research Center.

lent to $m_{\text{res}} J_5^a$ up to $O(a^2)$. Therefore when t is large enough such that low energy physics dominates, the ratio

$$R(t) = \frac{\langle \sum_{\vec{x}} J_{5q}^a(\vec{x}, t) \pi^a(0) \rangle}{\langle \sum_{\vec{x}} J_5^a(\vec{x}, t) \pi^a(0) \rangle} \quad (2)$$

should be equal to m_{res} . Fig. 1 shows that the plateau for $R(t)$ at $\beta = 6.0$, $L_s = 16$ starts from $t = 2$, and at $\beta = 5.7$, $L_s = 48$ it starts from $t = 4$.




Figure 1. $R(t)$ versus t at $\beta = 6.0$, $L_s = 16$ (circles) and $\beta = 5.7$, $L_s = 48$ (diamonds).

Being part of the mass term of the low energy effective Lagrangian close to the continuum, m_{res} should provide a universal description of the residual chiral symmetry breaking effects in all long distance observables. This is demonstrated in Figure 2 where m_{res} shows little m_f or volume dependence at $\beta = 5.70$, $L_s = 48$.

Figure 3 shows the L_s dependence of m_{res} at $\beta = 6.0$. The residual mass decreases with L_s , but the data points are poorly fit by a simple exponential. If we add a constant to the function, a much better fit can be obtained. This constant, which is the residual mass at infinite L_s , is about




Figure 2. The residual mass m_{res} versus m_f at $\beta = 5.70$, $L_s = 48$. The open circles are obtained from 106 configurations with volume $16^3 \times 32$, while the closed diamonds are obtained from 335 configurations with volume $8^3 \times 32$.

1 MeV. The five points can also be fit extremely well to a double-exponential function; this may suggest the existence of multiple decay modes. The exact asymptotic form for m_{res} as a function of L_s still needs further investigation. However for $L_s = 16$, m_{res} is already as low as 3.87(16) MeV in the $\overline{\text{MS}}$ scheme at 2 GeV, about 1/30 of the strange quark mass. This value is so small that the residual chiral symmetry breaking is not very important for these simulations. Even at the stronger coupling $\beta = 5.7$ ($a^{-1} \sim 1$ GeV), when L_s is increased from 32 to 48, m_{res} drops from 0.0105(2) to 0.00688(13), which is about 1/14 of the strange quark mass. Good chiral properties can be obtained by simulating at this value of L_s . Studies of this issue can also be found in [7,8].

3. CALCULATION OF f_π

As a test of these measurements and chiral properties, we calculate the pion decay constant using the pseudoscalar density correlator, which




Figure 3. The residual mass m_{res} versus L_s at $\beta = 6.0$. The dotted line is a simple exponential fit. The long dashed line is an exponential plus constant fit. The solid line is a double-exponential fit.

requires knowledge of the residual mass, and compare the results with those obtained from the axial vector current correlator.

Following the definition of the pion decay constant, f_π can be related to the amplitude of the axial current correlator A_{AA} :

$$A_{AA} = \left(\frac{f_\pi}{Z_A}\right)^2 \frac{m_\pi}{2}. \quad (3)$$

In Eq. 3, the renormalization factor Z_A is needed to relate the local axial current to the conserved current in the domain wall fermion formulation. Z_A can be measured by studying the large t behavior of the quantity $Z_A(t)$, defined as the ratio of the coupling between the conserved current and the pion to the coupling between the local current and the pion. Figure 4 shows that nice plateaus for $Z_A(t)$ can be found over the ranges $4 \leq t \leq 14$ and $18 \leq t \leq 28$ for both $\beta = 6.0$, $L_s = 16$ with $16^3 \times 32$ volume and $\beta = 5.7$, $L_s = 48$ with $8^3 \times 32$ volume. The L_s dependence of Z_A at $\beta = 6.0$ is presented in Fig. 5. For L_s ranging from 12

to 48, the change in Z_A is less than 1%. Therefore, the effect of finite L_s on the result for Z_A is negligible.




Figure 4. $Z_A(t)$ versus t at $\beta = 6.0$, $L_s = 16$ on $16^3 \times 32$ lattices (circles) and $\beta = 5.7$, $L_s = 48$ on $8^3 \times 32$ lattices (diamonds).

Another method to calculate f_π uses the relation between f_π and the amplitude of the pseudoscalar correlator A_{PP} :

$$A_{PP} = -\left(\frac{f_\pi}{m_f + m_{\text{res}}}\right)^2 \frac{m_\pi^3}{8}, \quad (4)$$

which directly depends upon the residual mass m_{res} .

Results for $\beta = 5.7$, $L_s = 48$ with lattice volume $8^3 \times 32$ show a discrepancy in the m_f dependence of f_π between these two methods. While the difference appears to become somewhat smaller at the larger volume of $16^3 \times 32$, we attribute the discrepancy between these two methods of calculating f_π to $O(a^2)$ errors.

Figure 6 shows the comparison of these two independent calculations at the weaker coupling $\beta = 6.0$. Consistent results are seen at each valence quark mass, and the f_π values measured by




Figure 5. Z_A versus L_s at $\beta = 6.0$ on $16^3 \times 32$ lattices. The dashed line fits the points to the form of $A + B \exp(-\alpha L_s)$.

extrapolation to $m_f = -m_{\text{res}}$ agree well with the physical value of 130 MeV, where m_ρ is used to set the scale. It is important to note that these good properties disappear if the value for m_{res} in Eq. 4 is replaced by 0 or by the x intercept of the simple linear fit to m_π^2 . Therefore, the good agreement seen between the two approaches serves as a consistency check of the residual mass analysis and a demonstration of the good chiral properties of the domain wall fermion formulation.

4. CONCLUSIONS

The chiral symmetry breaking effects in the domain wall fermion formulation can be characterized by a residual mass, which enters the effective quark mass in the low energy effective Lagrangian. This quantity can be determined accurately from the additional term in the divergence of axial current. Good chiral properties can be achieved for the quenched domain wall formulation for lattice spacing $a^{-1} \sim 1 - 2$ GeV. The results for m_{res} are further checked by our determinations of the pion decay constant.




Figure 6. $f_{\pi,K}$ versus m_f at $\beta = 6.0$ on $16^3 \times 32$ lattices from $\langle A_0^a(x)A_0^a(0) \rangle$ correlator ($\circ$) and $\langle \pi^a(x)\pi^a(0) \rangle$ correlator ($\diamond$). The solid symbols represent the extrapolations to $m_f = -m_{\text{res}}$ and interpolations to the kaon mass. The diamonds are slightly shifted to the right for clarity.

5. ACKNOWLEDGMENT

These calculations were performed on the QCDSF machines at Columbia and the RIKEN BNL Research Center.

REFERENCES

1. D. Kaplan, Phys. Lett. **B288** (1992) 342.
2. P. Chen *et. al.*, Nucl. Phys. **B** (Proc. Suppl.) **73** (1999) 204.
3. L. Wu (RIKEN-BNL-CU), Nucl. Phys. **B** (Proc. Suppl.) **83-84** (2000) 224.
4. T. Blum *et. al.*, hep-lat/0007038.
5. M. Wingate (RIKEN-BNL-CU), these proceedings, hep-lat/0009023.
6. V. Furman and Y. Shamir, Nucl. Phys. **B439** (1995) 54.
7. S. Aoki *et. al.*, Phys. Rev. Lett. **84**, (2000) 238, hep-lat/9904012.
8. A. A. Khan *et. al.*, hep-lat/0007014